The Large Hadron Collider Recreating the Big Bang

P. Gutierrez University of Oklahoma

April 9, 2009

Standard Model of FUNDAMENTAL PARTICLES AND INTERACTIONS

The Standard Model summarizes the current knowledge in Particle Physics. It is the guantum theory that includes the theory of strong interactions (guantum chromodynamics or QCD) and the unified theory of weak and electromagnetic interactions (electroweak). Gravity is included on this chart because it is one of the fundamental interactions even though not part of the "Standard Model."

FERMIONS

matter			
spin =	1/2,	3/2,	5/2,

Leptons spin = 1/2 Flavor Mass Electric			Quarl	Quarks spin = 1/2		
Flavor	Mass GeV/c ²	Electric charge	Flavor	Approx. Mass GeV/c ²	Electric charge	
ν_{e} electron neutrino	<1×10 ⁻⁸	0	U up	0.003	2/3	
e electron	0.000511	-1	d down	0.006	-1/3	
$ u_{\mu}^{\mu}$ muon neutrino	<0.0002	0	C charm	1.3	2/3	
$oldsymbol{\mu}$ muon	0.106	-1	S strange	0.1	-1/3	
$ u_{ au}^{ ext{ tau}}_{ ext{neutrino}}$	<0.02	0	t top	175	2/3	
au tau	1.7771	-1	b bottom	4.3	-1/3	

Spin is the intrinsic angular momentum of particles. Spin is given in units of ħ, which is the quantum unit of angular momentum, where $h = h/2\pi = 6.58 \times 10^{-25}$ GeV s = 1.05x10⁻³⁴ J s.

Electric charges are given in units of the proton's charge. In SI units the electric charge of the proton is 1.60×10^{-19} coulombs.

The energy unit of particle physics is the electronvolt (eV), the energy gained by one electron in crossing a potential difference of one volt. Masses are given in GeV/c² (remember $E = mc^2$), where 1 GeV = 10⁹ eV = 1.60×10⁻¹⁰ joule. The mass of the proton is 0.938 GeV/c² = 1.67×10⁻²⁷ kg.

Pro

Particles experiencing:

Particles mediating:

Strength relative to electromag 10⁻¹⁸ m

for two protons in nucleus

3×10⁻¹⁷ m

A neutron decays to a proton, an electron.

and an antineutrino via a virtual (mediating) W boson. This is neutron B decay

 $n \rightarrow p e^- \overline{\nu}_{o}$

for two u quarks at:

Baryons qqq and Antibaryons qqq Baryons are fermionic hadrons. There are about 120 types of baryons.					
Symbol	Name	Quark content	Electric charge	Mass GeV/c ²	Spin
р	proton	uud	1	0.938	1/2
p	anti- proton	ūūd	-1	0.938	1/2
n	neutron	udd	0	0.940	1/2
Λ	lambda	uds	0	1.116	1/2
Ω-	omega	SSS	-1	1.672	3/2

Matter and Antimatter

For every particle type there is a corresponding antiparticle type, denoted by a bar over the particle symbol (unless + or - charge is shown). Particle and antiparticle have identical mass and spin but opposite charges. Some electrically neutral bosons (e.g., Z^0 , γ , and $\eta_c = c\overline{c}$, but not $K^0 = d\bar{s}$) are their own antiparticles.

Figures

These diagrams are an artist's conception of physical processes. They are not exact and have no meaningful scale. Green shaded areas represent the cloud of gluons or the gluon field, and red lines the quark paths.



PROPERTIES OF THE INTERACTIONS

BOSONS



force carriers spin = 0, 1, 2, ...

nified Ele	ctroweak s	pin = 1	
lame	Mass GeV/c ²	Electric charge	
γ hoton	0	0	
W-	80.4	-1	c
W+	80.4	+1	C Ei TI
Z ⁰	91.187	0	T

Strong (color) spin = 1



olor Charge

ach quark carries one of three types of strong charge," also called "color charge." ese charges have nothing to do with the lors of visible light. There are eight possible types of color charge for gluons. Just as electri-

cally-charged particles interact by exchanging photons, in strong interactions color-charged particles interact by exchanging gluons. Leptons, photons, and W and Z bosons have no strong interactions and hence no color charge

Quarks Confined in Mesons and Baryons

One cannot isolate guarks and gluons: they are confined in color-neutral particles called hadrons. This confinement (binding) results from multiple exchanges of gluons among the color-charged constituents. As color-charged particles (quarks and gluons) move apart, the energy in the color-force field between them increases. This energy eventually is converted into additional quark-antiquark pairs (see figure below). The quarks and antiquarks then combine into hadrons; these are the particles seen to emerge. Two types of hadrons have been observed in nature: mesons $q\bar{q}$ and baryons qqq.

Residual Strong Interaction

Hadrons

Mesons

Not applicable

to quarks

20

The strong binding of color-neutral protons and neutrons to form nuclei is due to residual strong interactions between their color-charged constituents. It is similar to the residual electrical interaction that binds electrically neutral atoms to form molecules. It can also be viewed as the exchange of mesons between the hadrons.

Interaction	Gravitational	Weak	Electromagnetic	Stro	ong
	eravitational	(Electr	oweak)	Fundamental	Residual
Acts on:	Mass – Energy	Flavor	Electric Charge	Color Charge	See Residual Strong Interaction Note

Electrically charged

1

1

R

B⁰

Quarks, Gluons

Gluons

25

60

Not applicable

to hadrons

hadrons

70

Z⁰

 $p p \rightarrow Z^0 Z^0 + assorted hadrons$

Two protons colliding at high energy can produce various hadrons plus very high mass

particles such as Z bosons. Events such as this

one are rare but can yield vital clues to the

structure of matter

Ouarks, Leptons

 $W^+ W^- 7^0$

0.8

10-4

10-7

or

An electron and positron (antielectron) colliding at high energy can

nnihilate to produce B^0 and \overline{B}^0 mesons

via a virtual Z boson or a virtual photon

e⁺e⁻ → B⁰ B⁰

e⁺

e

All

Graviton

10-41

10-41

10-36

(not yet obse

e-

ve

	Mesons are bosonic hadrons. There are about 140 types of mesons.					
Symbol	Name	Quark content	Electric charge	Mass GeV/c ²	Spin	
π^+	pion	ud	+1	0.140	0	
К-	kaon	sū	-1	0.494	0	
ρ^+	rho	ud	+1	0.770	1	
B ⁰	B-zero	db	0	5.279	0	
η_{c}	eta-c	ςΣ	0	2 .980	0	

Mesons aa

The Particle Adventure

Visit the award-winning web feature The Particle Adventure at http://ParticleAdventure.org

This chart has been made possible by the generous support of: U.S. Department of Energy

U.S. National Science Foundation Lawrence Berkeley National Laboratory Stanford Linear Accelerator Center American Physical Society, Division of Particles and Fields BURLE INDUSTRIES INC.

©2000 Contemporary Physics Education Project. CPEP is a non-profit organiza-tion of teachers, physicists, and educators. Send mail to: CPEP, MS 50-308, Lawrence Berkeley National Laboratory, Berkeley, CA, 94720. For information on charts, text materials, hands-on classroom activities, and workshops, see:

http://CPEPweb.org



History of the Universe



What We Know







Fundamental Questions

- Standard Model
- Particles
- History
- Composition
- Questions
- ♦ LHC View
- LHC Diagram
- LHC Cavity
- LHC Magnet
- LHC Experiments
- ♦ LHC ATLAS
- ATLAS Cavern
- ✤ ATLAS
- Detector Objects
- Toroid
- Muon
- Pixel
- Pixel
- Pixel OU
- Higgs
- TDAQ
- The Grid
- Grid Locations

- Why do particles have mass (what is the origins of the Higgs mechanism)?
- Why more matter than anti-matter?
 - Are the forces unified?
 - Why three families of quarks and leptons?
 - Why is the mass of the top-quark so large?
 - Do quarks and leptons have a size or are they truly points?
- What is the dark matter composed of?
 - What is the nature of the dark energy?









Overall view of the LHC experiments.



























